


A1
1. [Amended] A method for controlling *Cryptosporidium parvum* in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound, such that *Cryptosporidium parvum* is controlled in said mammal, wherein said tetracycline compound inhibits more than 70% of *Cryptosporidium parvum* at a concentration less than 10 µg/ml.

2. [Amended] The method of claim 1, wherein said tetracycline compound is of formula I:



wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CHR^6 , S, NR^6 , or O;

R^2 , R^4 and $\text{R}^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, or heteroaromatic;

$\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen;

R^5 is hydroxy, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^6 , R^7 , R^8 and R^9 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

3. [Amended] The method of claim 2, wherein R^2 , $\text{R}^{2'}$, R^3 , R^{10} , R^{11} , and R^{12} are each hydrogen.

A2

45. [Amended] A method for treating a *Cryptosporidium parvum* related disorder in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound such that said mammal is treated for said disorder, wherein said tetracycline compound inhibits more than 70% of *Cryptosporidium parvum* at a concentration less than 10 µg/ml.

A3

-

(I)

and pharmaceutically acceptable salts thereof.


A3 cont'd
47. [Amended] The method of claim 46, wherein R^2 , $R^{2'}$, R^3 , R^{10} , R^{11} , and R^{12} are each hydrogen.

A4
58. [Amended] The method of claim 46, wherein said tetracycline compound is selected from the group consisting of 5-propionyl-6-cyclopentylsulfanylmethyl doxycycline; thiatetracycline; 9-cyclopent-1-enyl-doxycycline; 5-propionyl-9-tert-butyl-doxycycline; 9-tert-butyl doxycycline; 9-cyclohex-1-enylethynyl minocycline; and 6-cyclopentylsulfanylmethyl doxycycline.

A5
70. [Amended] The method of claim 46, wherein said supplementary agent is paromomycin.

Please add new claims 78-83:

A6
78. [New] A method for controlling *Cryptosporidium parvum* in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound, such that *Cryptosporidium parvum* is controlled in said mammal, wherein said tetracycline compound is of formula I:



wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CHR^6 , S, NR^6 , or O;

R^2 , R^4 and $\text{R}^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic (or a prodrug moiety);

$\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is alkanoyl;


R^6 , R^7 , R^8 and R^9 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

79. [New] A method for controlling *Cryptosporidium parvum* in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound, such that *Cryptosporidium parvum* is controlled in said mammal, wherein said tetracycline compound is of formula I:



wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$;

R^2 , R^4 and $\text{R}^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxy, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^6 , R^7 , R^8 and R^9 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;


R^{13} is hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

80. [New] A method for controlling *Cryptosporidium parvum* in a mammal, comprising administering to said mammal an effective amount of a tetracycline

compound, such that *Cryptosporidium parvum* is controlled in said mammal, wherein said tetracycline compound is of formula I:



wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CHR^6 , S, NR^6 , or O;

R^2 , R^4 and $\text{R}^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

R^2 , R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxy, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^6 , R^7 , and R^8 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;


R^9 is hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

81. [New] A method for treating a *Cryptosporidium parvum* related disorder in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound such that said mammal is treated for said disorder, wherein said tetracycline compound is of formula I:



(I)

wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$, CHR^6 , S, NR^6 , or O;

R^2 , R^4 and $\text{R}^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$\text{R}^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is alkanoyl;


R^6 , R^7 , R^8 and R^9 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

82. [New] A method for treating a *Cryptosporidium parvum* related disorder in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound such that said mammal is treated for said disorder, wherein said tetracycline compound is of formula I:



(I)

wherein:

X is $\text{CHC}(\text{R}^{13}\text{Y}'\text{Y})$;

Alc cont'd
 R^2 , R^4 and $R^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$R^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxy, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;


R^6 , R^7 , R^8 and R^9 are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R^{13} is hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.

83. [New] A method for treating a *Cryptosporidium parvum* related disorder in a mammal, comprising administering to said mammal an effective amount of a tetracycline compound such that said mammal is treated for said disorder, wherein said tetracycline compound is of formula I:



wherein:

X is $\text{CHC}(R^{13}Y'Y)$, CHR^6 , S , NR^6 , or O ;

R^2 , R^4 and $R^{4'}$ are each hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, arylalkyl, aryl, heterocyclic, heteroaromatic or a prodrug moiety;

$R^{2'}$, R^3 , R^{10} , R^{11} and R^{12} are each hydrogen or a pro-drug moiety;

R^5 is hydroxy, hydrogen, thiol, alkanoyl, aroyl, alkaroyl, aryl, heteroaromatic, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Al
confid
R⁶, R⁷, and R⁸ are each independently hydrogen, hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R⁹ is hydroxyl, halogen, thiol, alkyl, alkenyl, alkynyl, aryl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

R¹³ is hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

Y' and Y are each independently hydrogen, halogen, hydroxyl, cyano, sulfhydryl, amino, alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylamino, or an arylalkyl;

and pharmaceutically acceptable salts thereof.
